This product is designed specifically as a general non-acid cleaner and disinfectant for use in homes, hospitals, nursing homes, patient rooms, operating rooms, ICU areas, schools, food service establishments, transportation terminals, office buildings, manufacturing facilities, lodging establishments, retail businesses and athlete/recreational facilities where housekeeping is of prime importance in controlling the hazard of cross contamination. It is formulated to disinfect hard, non-porous, inanimate environmental surfaces: floors, walls, metal surfaces, stainless steel surfaces, glazed porcelain, glazed ceramic tile, plastic surfaces, bathrooms, shower stalls, bathtubs, and cabinets. For plastic and painted surfaces, spot test on an inconspicuous area before use. May be used in the kitchen on counters, sinks, appliances, and stove tops. A rinse with potable water is required for surfaces in direct contact with food. In addition, this product deodorizes those areas which generally are hard to keep fresh smelling, such as garbage storage areas, empty garbage bins and cans, basements, restrooms and other areas which are prone to odors caused by microorganisms, DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This product is not for use on critical or semi-critical medical device surfaces. Disinfection, deodorizing and cleaning -Remove gross filth or heavy soil prior to application of the product. Hold container six to eight inches from surface to be treated. Spray area until it is covered with the solution. Allow product to penetrate and remain wet for 3 minutes (for Bloodborne Pathogens; HIV-1, HBV, HCV, and HAV allow surface to remain wet for 10 minutes). No scrubbing is necessary. Wipe off with a clean cloth, mop or sponge. The product will not leave grit or soap scum. One-step cleaning and disinfection of toilet bowls and urinals: Remove water from bowl by forcing water over trap with a swab mop applicator. Press swab mop against side of the bowl to remove excess water from applicator. Apply 1 to 2 ounces of this product evenly, allowing sufficient time for product to cover entire surface of bowl for maximum cleaning. Swab bowl completely using a scrub brush or toilet mop, making sure to get under the rim. Let stand 3 minutes (for Bloodborne Pathogens: HIV-1. HBV. HCV. and HAV allow surface to remain wet for 10 minutes) and flush. To disinfect food service establishment food contact surfaces: countertops, appliances, tables, hold container six to eight inches from the surface to be treated and spray area until it is covered with the solution. Allow product to penetrate and remain wet for 3 minutes (for Bloodborne Pathogens: HIV-1, HBV, HCV, and HAV allow surface to remain wet for 10 minutes). Wipe off with a clean cloth, mop or sponge and thoroughly rinse all surfaces in the area with potable water. Heavily soiled areas must be pre-cleaned before using this product. Do not use on eating utensils, glassware and cookware.

Ŭsing approved AOAC test methods (under Good Laboratory Practices, [GLPs]), in the presence of 5% blood serum and a 3 minute contact time, unless otherwise noted, this product kills the following organisms on hard non-porous inanimate surfaces: BACTERICIDAL ACTIVITY: Pseudomonas aeruginosa, (ATCC 15442), Listeria monocytogenes, (ATCC 35152), Staphylococcus aureus, (ATCC 6589), Yersinia entercollitica, (ATCC 23715), Salmonella choleraesuis, (ATCC 10708), Enterococcus taecium, (ATCC 6569), Escherichia coli, (ATCC 11229), Corynebacterium ammoniagenes, (ATCC 6871), Escherichia coli O157:H7, (ATCC 43995), Salmonella tybii, (ATCC 6539)

EPA REG. NO. 1839-83-83106 EPA EST. NO. 3862-GA-1

Manufactured For:

### ALGONQUIN PRODUCTS CO.

1165 Allgood Road Suite 13 • Marietta, GA 30062 To Reorder Call: 866-578-4240

Fax: 866-578-4241 Part Number: 1036



# RTU GERM CLEANER

### **CLEANER / DEODORIZER**

- MILDEWSTAT (on hard inanimate surfaces)
   DISINFECTANT
- TUBERCULOCIDAL \*VIRUCIDE FUNGICIDE (against pathogenic fungi)
- MEETS OSHA BLOODBORNE PATHOGEN STANDARD FOR HIV. HBV. HCV. & HAV

#### ACTIVE INGREDIENTS

U.S. Patent No. 5,444,094

n-Alkyl (60% C14, 30% C16, 5% C12, 5% C18) dimethyl benzyl ammonium chlorides0.105	5%
n-Alkyl (68% C <sub>12</sub> , 32% C <sub>14</sub> ) dimethyl ethylbenzyl ammonium chlorides	5%
INERT INGREDIENTS 99.79	0%
Total	0%

Limpiador Desinfectante Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.(If you do not understand this label, find someone to explain it to you in detail.)

# KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID:
uct container or label with you when calling a poison control center or doctor, or going for treatment.
Hold eye open and rinse slowly and gently with water for 15-20 minutes.     Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.     Call a poison control center or doctor for treatment advice.
Call a Poison Control Center or doctor immediately for treatment advice.     Have person sip a glass of water if able to swallow.     Do not induce vomiting unless told to do so by the poison control center or doctor.     Do not give anything by mouth to an unconscious person.
Move person to fresh air.     if person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.     Call a poison control center or doctor for further treatment advice.

NOTE TO PHYSICIAN- Probable mucosal damage may contraindicate the use of gastric lavage

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

Net Contents: 1 Quart (946 ml)

ANTIBIOTIC-RESISTANT BACTERICIDAL ACTIVITY: Methicillin resistant Staphylococcus aureus (MRSA), (ATCC 3593), Vancomycin resistant Enterococcus faecalis (VRE), (ATCC 51575), Vancomycin intermediate resistant Staphylococcus aureus (VISA), (CDC Isolate 9921). This product exhibits effective disinfectant activity against Clostridium difficile vegetative form, ATCC 6899, when the treated surface is allowed to remain wet for 1 minute. TUBERCULOCIDAL ACTIVITY This product exhibits disinfectant efficacy against Mycobacterium tuberculosis (BCG) at 20 degrees Centigrade when the treated surface is allowed to remain wet for 5 minutes. "VIRUCIDAL ACTIVITY This product kills on hard, non-pronous inanimate surfaces when allowed to remain wet for a 10 minute contact time against: HIV-1 (associated with AIDS), Poliovirus Type 1, Hepatitis B Virus (HBV), Hepatitis C Virus (HCV), Canine Parvovirus, Hepatitis A Virus (HAV), Bovine Viral Darfmed Virus (BVDV).

This product, when used on environmental, inanimate hard surfaces exhibits effective virucidal activity against Norvalk virus and Feline Calicivirus, when the treated surface is allowed to remain wet for 30 seconds. This product, when used on environmental, inanimate hard surfaces exhibits virucidal activity against Human Coronavirus when the treated surface is allowed to remain wet for 20 minutes. This product, when used on environmental, inanimate hard surfaces exhibits virucidal activity against Rabies Virus when the treated surface is allowed to remain wet for 30 seconds. KILLS HIV-1, HBV, AND HCV, AND HAV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health care settings (Hospitals, Nursing Homes) or other settings in which there is an expected likelihood of solling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis CVirus (HZV), Hepatitis CVirus (HZV), and Hepatitis AVirus (HZV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1. HBV. HCV AND HAV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS. PERSONAL PROTECTION: When handling items soiled with blood or body fluids use disposable latex gloves, gowns, masks, or eve coverings, CLEANING PROCEDURES; Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product, CONTACT TIME: Allow surface to remain wet for 10 minutes, DISPOSAL OF INFECTIOUS MATERIALS: Blood and other body fluids should be autoclaved and disposed of according to local regulations for infectious waste disposal. MILDEWSTAT - To control mold and mildew on pre-cleaned, hard, non-porous surfaces spray surface to be treated making sure to wet completely. Let air dry. Repeat application at weekly intervals or when mildew growth appears. FUNGICIDAL ACTIVITY - This product is fungicidal against the pathogenic fungus, Trichophyton mentagrophytes (Athlete's Foot Fungus) when used with a 10 minute contact time on clean, nonporous, hard surfaces as found in bathrooms, shower stalls, locker rooms, or other areas commonly contacted by bare feet. EFFICACY TESTS HAVE DEMONSTRATED THAT THIS PRODUCT IS AN EFFECTIVE BACTERICIDE, FUNGICIDE, AND VIRUCIDE IN THE PRESENCE OF ORGANIC SOIL (5% BLOOD SERUM).

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

KEEP OUT OF REACH OF CHILDREN. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

#### STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL PESTICIDE STORAGE - Store in a dry place no lower in temperature than 50°F or higher than 120°F. CONTAINER DISPOSAL - Pastic Containers: Tiple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. For containers 1 gallon or less: Do not reuse empty containers (bottle, can, bucket). Wrap container and put in trash.